

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CCL26/Eotaxin-3 in direct ELISAs and Western blots.
Source	Monoclonal Rabbit IgG Clone # 2152H
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human CCL26/Eotaxin-3 Thr24-Leu94 Accession # Q9Y258
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Eotaxin-3, also named CCL26 or SCYA26, is a novel human CC chemokine that maps to chromosome 7q11.2, within 40 kilobases of the Eotaxin-2 loci. Eotaxin-3/CCL26 has been shown to be constitutively expressed in the heart and ovary. In addition, low levels of Eotaxin-3/CCL26 expression can also be detected in various tissues. The expression of Eotaxin-3/CCL26 in vascular endothelial cells has been shown to be up-regulated by IL-13 and IL-4.

Eotaxin-3/CCL26 cDNA encodes a 94 amino acid (aa) residue protein with a putative signal peptide of either 23 or 26 aa residues. Recombinant Eotaxin-3/CCL26 has been produced in insect cells using a baculovirus expression system and shown to contain 71 aa residues. Recombinant Eotaxin-3/CCL26 is chemotactic for eosinophils and PHA-activated T cells. Eotaxin-3/CCL26 induces calcium flux in eosinophils as well as in CCR3-transfected cells. Eotaxin-3/CCL26 has also been shown to cross-desensitize cells to other CCR3 ligands. Both the 71 aa residue and 68 aa residue variants of recombinant Eotaxin-3 have been expressed in *E. coli* and found to have equal potency in inducing chemotaxis of a human CCR3-transfected cell line.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.