

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Desmoglein-4 in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human Desmoglein-1, -2, or -3 is observed and approximately 40% cross-reactivity with recombinant mouse Desmoglein-4 is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 683135
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Desmoglein-4 Glu24-Ala631 Accession # Q86SJ6
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Desmoglein-4 (DSG4) is a 110-130 kDa type I transmembrane glycoprotein in the cadherin family of calcium-dependent cell adhesion molecules. Desmogleins interact with transmembrane Desmocollins to form the adhesive components of desmosomes on epithelial cells. Desmoglein-4 contains four cadherin repeats in its extracellular region and two desmoglein repeat domains in its cytoplasmic region. Alternate splicing generates an additional isoform with a substitution in the cytoplasmic region. Desmoglein-4 is expressed on keratinocytes and trichocytes in the epidermis and hair follicles. It is required for hair follicle development, and DSG4 mutations are associated with autosomal recessive hypotrichosis. Within amino acids 24-631 (which includes the 26 aa propeptide and the ECD), human Desmoglein-4 shares 82% aa sequence identity with mouse and rat Desmoglein-4.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.