

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Activin B β B subunit in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human (rh) Inhibin B or rhActivin A is observed. In Western blots, approximately 20% cross-reactivity with rhInhibin B
Source	Monoclonal Mouse IgG ₁ Clone # 146807
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Activin B Gly293-Ala407 Accession # Q53T31
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Activins and inhibins, members of the TGF- β superfamily, are disulfide-linked dimeric proteins that were originally purified from gonadal fluids as proteins that stimulated or inhibited, respectively, pituitary follicle stimulating hormone (FSH) release. These proteins have since been shown to have a wide range of biological activities including: mesoderm induction, neural cell differentiation, bone remodeling, hematopoiesis and reproductive physiology. Activins/inhibins are produced as precursor proteins with an amino-terminal propeptide that is cleaved to release the carboxy-terminal bioactive ligands. Activins are homodimers or heterodimers of the various β subunit isoforms, while inhibins are heterodimers of a unique α subunit and one of the various β subunits. Five β subunits (mammalian β_A , β_B , β_C , β_E and *Xenopus* β_D) have been cloned. The activin/inhibin nomenclature reflects the subunit composition of the proteins: activin A (β_A - β_A), activin B (β_B - β_B), activin AB (β_A - β_B), inhibin A (α - β_A) and inhibin B (α - β_B). At present, little is known about the contribution of the other β subunits to activin or inhibin formation and biology. At the amino acid sequence level, the mature human β_B subunit is greater than 98% identical to mouse β_B , while the human and mouse α subunits share approximately 80% identity. Similarly to other TGF- β family members, activins exert their biological activities through binding to the heterodimeric complex composed of two membrane spanning serine-threonine kinases designated as type I and type II. Two forms of activin receptor type I (Act RI-A and Act RI-B) and two forms of activin receptor type II (Act RII-A and Act RII-B) have been identified. Activin binds directly to Act RII, the complex then associates with Act RI and initiates signaling. Besides activins, Act RII has been shown to bind certain other TGF- β superfamily members. Inhibin A has been shown to bind with low-affinity to Act RII. The existence of a distinct inhibin-specific receptor and/or signal transduction pathway has been hypothesized.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.