

DESCRIPTION


| | |
|---------------------------|--|
| Species Reactivity | Human |
| Specificity | Detects human Fcε RIα in direct ELISAs. |
| Source | Monoclonal Mouse IgG _{2B} Clone # 773704 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | Mouse myeloma cell line NS0-derived recombinant human Fcε RIα Val26-Gln205 Accession # P12319 |
| Conjugate | PerCP (Peridinin-chlorophyll Protein Complex) Excitation Wavelength: 482 and 564 nm Emission Wavelength: 675 nm |
| Formulation | Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|-----------------------------|-----------|
| Flow Cytometry | 10 μL/10 ⁶ cells | See Below |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

The α subunit of the high affinity IgE receptor (Fcε RIα or Fcε RIA) is a 45-48 kDa IgE-binding type I transmembrane glycoprotein of the multichain immune recognition (MIRR) family (1, 2). The receptor, Fcε RI, is a tetrameric complex of one α, one β and two disulfide-linked γ subunits (αβγ₂) on mast cells and basophils (1). An alternate trimeric form (αγ₂) is expressed on human, but not rodent, mast cells, basophils, eosinophils, visceral smooth muscle, platelets, intestinal columnar epithelium and professional antigen presenting cells (3). While the γ subunit is essential for expression of Fcε RIα on the cell surface and for cell signaling, the β subunit, when present, increases the half-life of the Fcε RI complex on the cell surface (3, 4). An isoform of the β subunit, βT, blocks processing of the α subunit and its cell surface expression (2, 3, 5). Human Fcε RIα cDNA encodes 257 amino acids (aa) including a 25 aa signal sequence, a 180 aa extracellular domain that contains two Ig-like domains that bind IgE plus an endoplasmic reticulum retention motif, a 21 aa transmembrane sequence and a 32 aa cytoplasmic domain (1, 3, 6). Over aa 26-205, human Fcε RIα shares 52% aa sequence identity with mouse Fcε RIα. Binding of IgE alone increases surface expression of Fcε RI, while crosslinking of IgE/Fcε RI complexes by IgE ligands (allergens) initiates receptor internalization and signaling (2, 4, 5). Mast cell and basophil activation by IgE/Fcε RI crosslinking causes degranulation, resulting in the release of histamine, leukotrienes, prostaglandins, and other mediators of immediate-type and late-phase allergic reactions. Circulating autoantibodies that crosslink Fcε RIα are often found in patients with chronic urticaria (7). Fcε RI on human antigen presenting cells mediates uptake and processing of allergens for presentation by class II MHC (2, 3). Fcε RI expression on human DC and Langerhans cells is up-regulated during allergic reactions (atopy) and correlates with serum IgE concentration (3).

References:

1. Shimizu, A. *et al.* (1988) Proc. Natl. Acad. Sci. USA **85**:1907.
2. Abramson, J. and I. Pecht (2007) Immunol. Rev. **217**:231.
3. Sutton, B.J. and A.M. Davies (2015) Immunol. Rev. **268**:222.
4. Yamasaki, S. and T. Saito (2008) J. Pharmacol. Sci. **106**:336.
5. Brenzovich, J. *et al.* (2009) J. Leukoc. Biol. **86**:1351.
6. Cauvi, D.M. *et al.* (2006) J. Biol. Chem. **281**:10448.
7. Kikuchi, Y. *et al.* (2001) J. Allergy Clin. Immunol. **107**:1056.