

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human NK1R in ELISAs.
Source	Monoclonal Mouse IgG ₃ Clone # 694501
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	HEK293 human embryonic kidney cell line transfected with human NK1R Accession # P25103
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.

*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 µL/10 ⁶ cells	See Below

DATA

Flow Cytometry



Detection of NK1R in HEK293 Human Cell Line Transfected with Human NK1R and eGFP by Flow Cytometry HEK293 human embryonic kidney cell line transfected with (A) human NK1R or (B) irrelevant protein, and eGFP was stained with Mouse Anti-Human NK1R PE-conjugated Monoclonal Antibody (Catalog # FAB66871P) or Mouse IgG₃ PE-conjugated Isotype Control (Catalog # IC007P, data not shown). View our protocol for [Staining Membrane-associated Proteins](#).

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

NK1R (Neurokinin-1 receptor), gene name TACR1 (tachykinin receptor-1), is an ~80 kDa 7-transmembrane glycoprotein receptor for the proinflammatory tachykinin neuropeptide, substance P. NK1R is constitutively or inducibly expressed on a wide variety of cells, including monocytes, macrophages, microglia, lymphocytes, neutrophils, mast cells, and neurons. The 407 amino acid (aa) human NK1R contains 89 extracellular aa over multiple segments which collectively share 92% aa identity with corresponding regions of mouse and rat NK1R. A ~50 kDa short isoform that ends at aa 311 lacks almost all of the C-terminal cytoplasmic signaling region. It is the only form expressed in human monocytes and undifferentiated THP-1 cells, and its function appears to differ from the long isoform.