

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Protocadherin-15 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 740523
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Protocadherin-15 Tyr28-Ala1376 Accession # Q96QU1
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Protocadherin-15 (PCDH-15) is a 250-270 kDa "other" member of the nonclustered group of the protocadherin family of molecules. In the adult, it is expressed on the base of hair cell stereocilia in the Organ of Corti, on cerebellar granule cells, and on rod and cone photoreceptors. In the ear, it is expressed as a homodimer, and binds (in trans) to cadherin 23 homodimers on adjacent stereocilia. Mature human PCDH-15 is a type I transmembrane protein that is 1929 amino acids (aa) in length. It contains a 1350 aa extracellular domain (ECD) (aa 27-1376) plus a 558 aa cytoplasmic region. There are 11 cadherin domains in the ECD (aa 40-1259). Multiple splice forms are reported. One is a soluble 110 kDa form that shows a five aa substitution for aa 957-1955. Another contains a seven aa substitution for aa 1119-1125. Others show substitutions or deletions in either the extracellular or cytoplasmic domains. Over aa 27-1376 (the ECD), human PCDH-15 shares 95% aa sequence identity with mouse PCDH-15.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.