

DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human GPR12 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 584940
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with human GPR12 Accession # P47775
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.

APPLICATIONS			
Please Note: Optimal dilutions should be	e determined by each laboratory for each applicatio	n. General Protocols are available in the Technical Information section on our website.	
	Recommended Concentration	Sample	
Flow Cytometry	0.25 μg/10 ⁶ cells	See Below	

DATA	
Flow Cytometry	Detection of GPR12 in HEK293 Human Cell Line Transfected with Human GPR12 and eGFP by Flow Cytometry. HEK293 human embryonic kidney cell line transfected with (A) human GPR12 or (B) irrelevant transfectants and eGFP was stained with Mouse Anti-Human GPR12 PE-conjugated Monoclonal Antibody (Catalog # FAB6738P). Quadrant markers were set based on control antibody staining (Catalog # IC002P). View our protocol for Staining Membrane-associated Proteins.
eGFP	
PREPARATION AND S	TORAGE
Shipping	The product is shipped with polar packs. Upon receipt, sto
Stability & Storage	 Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as suppl

BACKGROUND

GPR12 is a 7-transmembrane receptor for sphingosylphosphorylcholine. Human GPR12 and the related GPR3 and GPR6 are all highly expressed on neurons of the central nervous system, while GPR12 is particularly expressed in the limbic system. GPR12 engagement promotes proliferation and maturation of postmitotic neurons through an inhibitory G protein and cAMP signaling pathway. Human GPR12 cDNA encodes 334 amino acids (aa), 89 of which are extracellular; these share 86% aa identity with mouse and rat GPR12.

Rev. 12/5/2018 Page 1 of 1



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449