

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FCRL6/FcRH6 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) FCRL1, rhFCRL2, rhFCRL3, rhFCRL4, rhFCRL5, rhFcRN, or rhFCRLB/FCRY is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 672505
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human FCRL6/FcRH6 Lys16-Leu312 Accession # Q6DN72
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	See Below

DATA

Flow Cytometry



Detection of FCRL6/FcRH6 in Human Peripheral Blood Lymphocytes by Flow Cytometry. Human peripheral blood lymphocytes were stained with Mouse Anti-Human NCAM-1/CD56 APC-conjugated Monoclonal Antibody (Catalog # [FAB2408A](#)) and either (A) Mouse Anti-Human FCRL6/FcRH6 Alexa Fluor® 488-conjugated Monoclonal Antibody (Catalog # [FAB6745G](#)) or (B) Mouse IgG₁ Alexa Fluor 488 Isotype Control (Catalog # [IC002G](#)). View our protocol for [Staining Membrane-associated Proteins](#).

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

FCRL6 (Fc Receptor-Like Protein 6), also known as FcRH6 and IFGP6, is a 66 kDa member of the Ig Superfamily. It is found on CD56dim CD16⁺ NK cells, CD56⁺ CD3⁺ NKT cells, gd T cells, and effector plus effector-memory CD8⁺ T cells. FCRL6 is not a receptor for immunoglobulin. It does, however, bind to isoforms of HLA-DR that are expressed on APCs. Given that FCRL6 contains a cytoplasmic ITIM, FCRL6 may function as an inhibitory receptor for MHC Class II. Mature human FCRL6 is a 415 amino acid (aa) type I transmembrane protein. It contains a 288 aa extracellular region (aa 20-307) that shows three C2-type Ig-like domains (aa 20-293), and a 106 aa cytoplasmic domain. There are multiple potential splice variants. One shows a deletion of aa 105-200 accompanied by a 13 aa substitution for aa 394-434; a second possesses a 15 aa substitution for aa 383-434; and a third shows an alternative start site seven aa upstream of the standard site accompanied by a 13 aa substitution for aa 394-434. Over aa 16-312, human FCRL6 shares only 33% aa sequence identity with the mouse counterpart to FCRL6.

Human FCRL6/FcRH6 Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 672505

Catalog Number: FAB6745G

100 µg

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.