

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human LILRA5/CD85f in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant mouse LILRC1, recombinant rat LILRC1, recombinant human (rh) ILT7/LILRA4, rhILT4, rhILT5, or rhLILRB4/LIR5 is observed. In Western blots,
Source	Monoclonal Mouse IgG _{2A} Clone # 711827
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human LILRA5/CD85f Gly42-Arg268 Accession # A6NI73
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

ILT11 (also leukocyte immunoglobulin-like receptor 9 (LIR-9), CD85f, and LILRA5) is a 39-40 kDa Group 2 member of the LILR family of innate immune receptors. It is expressed on monocytes and perhaps neutrophils, and its activation results in the secretion of proinflammatory cytokines such as TNF-α and IL-1β. Mature human ILT11 is a 258 amino acid (aa) type I transmembrane glycoprotein. It has a 227 aa extracellular domain (aa 42-268) that contains two C2-type Ig-like domains (aa 51-136 and 142-230), and a 10 aa cytoplasmic tail. ILT11 has three potential splice forms. One is a 35 kDa soluble form of the molecule described above that shows a 27 aa substitution for aa 239-299. The other two splice forms are analogous to the above membrane and soluble forms, but demonstrate signal sequence cleavage further downstream after Ala51. Over aa 41-268, human ILT11 shares 58% aa identity with mouse ILT11.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.