Human IL-1 RAcP/IL-1 R3 Alexa Fluor® 700-conjugated Antibody

Monoclonal Mouse IgG_1 Clone # 89412

Catalog Number: FAB676N
100 Tests

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-1RAcP/IL-1 R3 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) IL-1 RI, rhIL-1 RII, rhST2, rhIL-18 Rα, IL-1 Rrp2, rhIL-18 RAP, rhSIGIRR, rhIL-1 RAPL1, rhIL-1 RAPL2, rhTLR1, rhTLR2, rhTLR3, rhTLR4, rhTLR5, rhTLR7, rhTLR8, rhTLR9, rhTLR10, rhMD-1, rhMD-2, or recombinant mouse RP105 is observed.
Source	Monoclonal Mouse IgG_1 Clone # 89412
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	*S. frugiperda* insect ovarian cell line *Sf*21-derived recombinant human IL-1RAcP/IL-1 R3 Ser21-Glu359 (predicted) Accession # Q9NPH3
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: *Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.*

	Recommended Concentration	Sample
Flow Cytometry	5 µL/10^6 cells	See Below

DATA

Flow Cytometry



Detection of IL-1 RAcP/IL-1 R3 in Human Monocytes by Flow Cytometry. Human peripheral blood monocytes were stained with Mouse Anti-Human IL-1 RAcP/IL-1 R3 Alexa Fluor® 700-conjugated Monoclonal Antibody (Catalog # FAB676N, filled histogram) or isotype control antibody (Catalog # IC002N, open histogram). View our protocol for Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	**Protect from light. Do not freeze.** ● 12 months from date of receipt, 2 to 8 °C as supplied.

bio-techne®
Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956
USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373
Europe | Middle East | Africa TEL +44 (0)1235 529449

Human IL-1 RAcP/IL-1 R3
Alexa Fluor® 700-conjugated Antibody

Monoclonal Mouse IgG_1 Clone # 89412

Catalog Number: FAB676N
100 Tests

BACKGROUND

IL-1 Receptor Accessory Protein (also IL-1 R3) is a ubiquitous 70-90 kDa member of the interleukin-1 receptor family of proteins (1-5). It serves as a non-ligand-binding accessory component of the receptors for IL-1α, IL-1β, and IL-33 (6, 7). Together with IRAK4 and MyD88, it generates a functional signaling complex with IL-1 RI; by itself, it generates a non-signaling, but high-affinity binding complex with IL-1 RII (8). In addition, it interacts with ST2 on mast cells and Th2 T cells to create a functional IL-33 receptor complex (7). Mature human IL-1 RAcP is a type I transmembrane glycoprotein that is 550 amino acids in length. It contains a 347 amino acid (aa) extracellular region (aa 21-367), a 21 aa transmembrane segment, and a 182 aa cytoplasmic domain (9). The extracellular region shows three C2-type Ig-like domains, the most membrane proximal of which is suggested to be responsible for dimerization with IL-1 RI (10). There are three alternative splice forms reported for IL-1 RAcP. One is transmembrane and shows a 239 aa substitution for the C-terminal 122 amino acids (11). The other two are soluble; one shows a six aa substitution for aa 351-570, while a second shows a 45 aa substitution for aa 302-579 (12, 13). The soluble receptor isoforms appear to be inhibitory to IL-1 signaling. When present with soluble IL-1 RII, soluble IL-1 RAcP increases the IL-1 binding affinity of IL-1 RII more than 100-fold, thus neutralizing the effects of IL-1 (14). The human and mouse IL-1 RAcP precursors are 89% aa identical; within the extracellular region, they share 86% aa identity.

References:

1. Subramaniam, S. *et al.* (2004) Dev. Comp. Immunol. **28**:415.
2. Boraschi, D. and A. Tagliabue (2006) Vitam. Horm. **74**:229.
3. Dunne, A. and L.A.J. O'Neill (2003) Sci STKE. Feb 25;2003(171):re3.
4. Huang, J. *et al.* (1997) Proc. Natl. Acad. Sci. USA **94**:12829.
5. Greenfeder, S. A. *et al.* (1995) J. Biol. Chem. **270**:13757.
6. Brikos, C. *et al.* (2007) Mol. Cell. Proteomics **6**:1551.
7. Chackerian, A.A. *et al.* (2007) J. Immunol. **179**:2551.
8. Lang, D. *et al.* (1998) J. Immunol. **161**:6871.
9. SwissProt. Accession # Q9NPH3.
10. Yoon, D-Y. and C.A. Dinarello (1998) J. Immunol. **160**:3170.
11. Lu, H-L. *et al.* (2008) Mol. Immunol. **45**:1374.
12. Jensen, L.E. *et al.* (2000) J. Immunol. **164**:5277.
13. Jensen, L.E. and A.S. Whitehead (2003) Cell. Signal. **15**:793.
14. Smith, D.E. *et al.* (2003) Immunity **18**:87.

PRODUCT SPECIFIC NOTICES

bio-techne®
Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956
USA TEL 800 343 7475 **Canada** TEL 855 668 8722 **China** TEL +86 (21) 52380373
Europe | Middle East | Africa TEL +44 (0)1235 529449