

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human NLRP1/NALP1 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 447916
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human NLRP1/NALP1 Gly1331-Leu1429 Accession # Q9C000
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

NALP1 (Nacht, Leucine-rich repeat and PYD domain containing protein 1; also NAC, CARD7, DEFCAP and CLR17.1) is a 160-170 kDa member of the NLRP family of molecules. It is expressed in dendritic cells, neutrophils, T and B cells, alveolar macrophages, permatogonia, neurons, plus intestinal columnar epithelium, and is found in both cytoplasm and nucleus. NALP1 promotes apoptosis plus IL-1 β and IL-18 maturation by activating caspase-1 and -5. It does so by forming a 700 kDa inflammasome comprised of NALP1, ASC plus caspase-1 and -5. Bacterial wall peptidoglycan binds to NALP1, promoting ATP binding, NALP1 oligomerization, and caspase activation. Human NALP1 is 1473 amino acids (aa) in length. It contains an N-terminal DAPIN domain (aa 1-92), a NACHT domain (aa 328-637), seven consecutive LRRs (aa 704-1236) and one CARD region (aa 1374-1463). Alternate splice forms exist that range in size from 70 kDa-150 kDa. Either individually, or in combination, there can be a 19 aa substitution for either aa 1353-1473 or aa 1368-1472, a deletion of aa 91-260, 958-987 or 1262-1305, and a 43 aa insert after Leu785. Over aa 1-323, human NALP1 shares less than 20% aa identity with mouse NALP1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.