

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human sFRP-2 in direct ELISAs and Western blots. In direct ELISAs, 100% cross-reactivity with recombinant mouse (rm) sFRP-2 is observed and no cross-reactivity with recombinant human (rh) sFRP-1 or rh sFRP-5 is observed. In Western blots, no cross
Source	Monoclonal Mouse IgG _{2B} Clone # 755644
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human sFRP-2 Leu25-Cys295 Accession # Q96HF1
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Secreted Frizzled Related Protein-2 (sFRP-2) belongs to a family of Wnt-binding proteins with homology to the ligand-binding domain of the Frizzled transmembrane Wnt receptors. The sFRP proteins are approximately 30-35 kDa in size and contain an N-terminal Frizzled-like domain with 10 conserved cysteines and a Netrin-like C-terminal domain (1, 2). Mature human sFRP-2, also known as SARP-1, SDF-5, and FRP-2, shares 99% aa sequence identity with mouse and rat sFRP-2 (3). sFRP-2 is widely expressed during embryogenesis and in the adult in tissues including the eye, heart, lung, colon, intestine, smooth muscle, pancreas, prostate, testis, kidney, brain, teeth and joints, craniofacial mesenchyme, and preadipocytes (3-6). Depending on the context, sFRP-2 can exert either positive or negative effects on Wnt signaling (7-9). It also inhibits BMP-induced effects (8, 10). sFRP-2 can be incorporated into the extracellular matrix through interactions with Fibronectin and Integrin α5β1 (11). sFRP-2 plays a variety of roles during tissue morphogenesis including inhibition of the planar cell polarity pathway and myoblast and osteoblast differentiation (8, 10, 12, 13). sFRP-2 is also expressed in multiple myeloma and glioma in which it promotes tumorigenicity (10, 14). At physiological concentrations sFRP-2 enhances BMP-1 mediated proteolysis of Pro-Collagen I, whereas at higher concentrations it inhibits BMP-1 activity (15, 16). This difference is significant considering that sFRP-2 is up-regulated in fibrotic areas of the heart following myocardial infarction (15, 16). Elevated levels of sFRP-2 then promote the recovery of cardiac function by reducing collagen deposition, remodeling, and calcification and by promoting the engraftment of mesenchymal stem cells into the heart (8, 15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.