

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human PAK2 in direct ELISAs. In direct ELISAs, approximately 5% cross-reactivity with recombinant human (rh) PAK1 and no cross-reactivity with rhPAK1B, 3, 4, 6, or 7 is observed.
Source	Monoclonal Mouse IgG ₃ Clone # 677411
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human PAK2 Leu127-Gln221 Accession # Q13177
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

PAK2 (p21-activated kinase-2) is a widely expressed 58-62 kDa cytoplasmic ser/thr kinase of the PAK family, with substrates including ribosomal S6 and histone H4 proteins. The 524 amino acid (aa) human PAK2 contains phosphorylation, acetylation (K38, K128) and myristoylation (G213) sites, a CRIB (cdc42/Rac interactive binding) domain (aa 74-87), a caspase-3 cleavage site after aa 212, and a kinase domain (aa 249-499). Full-length PAK2 stimulates cell growth and motility, while the caspase cleaved PAK2 kinase domain (34 kDa) is involved in cell death. Within the region used as an immunogen, human PAK2 shares 87% aa identity with mouse and rat PAK2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.