Human GITR/TNFRSF18 Fluorescein-conjugated Antibody

Monoclonal Mouse IgG_1 Clone # 110416

Catalog Number: FAB689F

100 TESTS, 25 TESTS

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human GITR in direct ELISAs and Western blots. Does not cross-react with recombinant human (rh) 4−1BB, recombinant mouse (rm) 4−1BB, rhCD27, rmCD27, rhCD30, rmCD30, rhCD40, rmCD40, rhDR3, rhDR6, rhEDAR, rmEDAR, rhFas, rmFAS, rmGITR, rhHVEM, rhLymphotoxin Rβ, rmLymphotoxin Rβ, rhNGF R, rhOPG, rmOPG, rhRANK, rmRANK, rhTAJ, rhTNF RI, rmTNF RI, rhTNF RII, or rmTNF RII.
Source	Monoclonal Mouse IgG_1 Clone # 110416
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human GITR/TNFRSF18 Gln26-Glu161 (Thr43Ala) Accession # Q9Y5U5
Conjugate	Fluorescein Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm (FITC)
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: *Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.*

	Recommended Concentration	Sample
Flow Cytometry	10 µL/10^6 cells	See Below

DATA

Flow Cytometry



Detection of GITR/TNFRSF18 in Human PBMCs by Flow Cytometry. Human peripheral blood mononuclear cells (PBMCs) treated with 1 µg/mL PHA for 24 hours were stained with Mouse Anti-Human GITR/TNFRSF18 Fluorescein-conjugated Monoclonal Antibody (Catalog # FAB689F, filled histogram) or isotype control antibody (Catalog # IC002F, open histogram). View our protocol for Staining Membrane-associated Proteins.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	**Protect from light. Do not freeze.** ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

GITR (glucocorticoid-induced tumor necrosis factor receptor, also named AITR, activation-inducible TNF R family member), is a 228 amino acid (aa) type I transmembrane protein belonging to the TNF R family and has been designated TNFRSF18. The GITR cytoplasmic domain has striking homology with the cytoplasmic domain of 4-1BB and CD27. Human GITR shares 55% homology with murine GITR. GITR is expressed at low levels in peripheral blood T cells, bone marrow, thymus, spleen, and lymph nodes. In contrast to mouse GITR, expression of human GITR is not induced by treatment with dexamethasone, but is up−regulated by antigen stimulation or by treatment with anti-CD3 plus anti-CD28, or PMA plus ionomycin. Human GITR ligand was identified from human umbilical vein endothelial cells and is a 177 aa polypeptide belonging to the TNF superfamily (TNFSF18). Ligation of GITR can activate NF-κB through TRAF2, and protect T cells from TCR activation-induced cell death. It has been proposed that GITR ligand and GITR may modulate T lymphocyte functions.

References:

1. Nocentini, G. *et al.* (1997) Proc. Natl. Acad. Sci. USA **94**:6216.
2. Kwon, B. *et al.* (1999) J. Biol. Chem. **274**:6056.
3. Gurney, A.L. *et al.* (1999) Current Biology **9**:215.
4. Kwon, B. *et al.* (1999) Current Opinion in Immunology **11**:340.

bio-techne
bio-techne.com
Global info@bio-techne.com techsupport@bio-techne.com
North America TEL 800 343 7475 Europe | Middle East | Africa TEL +44 (0)1235 529449
China info.cn@bio-techne.com TEL +86 (21) 52380373
Rest of World bio-techne.com/find-us/distributors TEL +1 612 379 2956