

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CD11c in direct ELISAs and Western blots. In direct ELISAs, 50%-100% cross-reactivity with recombinant human (rh) Integrin α X β 2 and recombinant mouse (rm) Integrin α L is observed, 10%-25% cross-reactivity with rhIntegrin
Source	Monoclonal Rat IgG _{2B} Clone # 435421
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse CD11c Phe20-Pro1116 Accession # Q9QXH4
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CD11c, also known as the Integrin α X subunit, is a 150 kDa type I transmembrane protein that noncovalently heterodimerizes with the β 2 subunit (CD18) to form α X/ β 2, also known as p150/p95 and complement receptor type 4 (CR4). Integrin α X β 2 is expressed on macrophages, dendritic cells, hairy cell leukemias and some other leukocyte subsets. The 1097 aa mouse CD11c extracellular domain shares 71% and 87% amino acid (aa) identity with human and rat CD11c, respectively. One potential α X isoform is truncated at aa 828. Some adhesion partners of α X β 2 are shared with α M β 2/CD11b/CD18 (Complement iC3b, ICAMs, vWF and Fibrinogen) while others (Osteopontin, Thy-1, Plasminogen, Heparin) are unique. Unlike α M β 2, it is not constitutively active. α X β 2 adhesion mediates proliferation, degranulation, chemotactic migration, and phagocytosis of complement-opsonized particles.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.