

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Tenascin XB2 in ELISAs and Western blots. Also detects Tenascin X in Western blots. In direct ELISAs, no cross-reactivity with recombinant human Tenascin R is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 766842
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Tenascin XB2 Met1-Tyr673 Accession # AAH33740
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Tenascin XB2 (TN-XB2; also TN-XB-Short/S and tenascin-X isoform 2) is a 73-75 kDa member of the tenascin family of extracellular matrix molecules. It is expressed by adrenal gland, and represents the use of an intronic promoter that is tied to an internal start site within the TN-XB gene. A duplication of the TN-XB2 region has created a gene called TN-XA that codes for the same ORF but is not translated. Although TN-XB2 has been reported to bind to tropoelastin, plus collagens I, III and V, other studies using soluble fragments of TN-XB report only tropoelastin interaction. Human TN-XB2 is 673 amino acids (aa) in length. It contains four fibronectin type III repeats (aa 82-442), and one C-terminal fibrinogen-like globular region (aa 449-664). Antibodies to TN-XB2 recognize full-length 465 kDa Tenascin X (TN-XB), 75 kDa and 140-150 kDa C-terminal fragments. Although human TN-XB2 shares 84% aa identity with full-length mouse TN-XB in its C-terminus, there does not appear to be an equivalent alternative start site in the mouse gene.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.