

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human SARM1 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 754711
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human SARM1 Lys556-Thr724 Accession # Q6SZW1
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SARM1 (Sterile alpha and TIR motif-containing protein 1; also SAM domain-containing protein 2 and TIR-1 homolog) is a 73-75 kDa member of the TLR adaptor family of molecules. It is both nuclear and cytoplasmic, and expressed in monocytes, neurons, and retinal photoreceptor cells, where it demonstrates disparate activities. In monocytes, SARM1 is a specific inhibitor of TRIF-dependent TLR-3 and -4 signaling, and appears to block MAPK phosphorylation. In neurons, SARM1 regulates microtubule stability, and thus axon and dendrite elongation. And in retinal photoreceptor cells, SARM1 complexes with Na/K ATPase to create a cell surface receptor for retinoschisin. Human SARM1 is 724 amino acids (aa) in length. It contains an N-terminal polybasic motif (aa 1-20) and a Gly-rich region (aa 22-90) that may anchor the molecule to intracellular membranes. It also possesses two SAM domains (aa 412-548) plus a TIR region (aa 559-657) that appears to interact with TIRF. Proteolytic cleavage products of 35 kDa and 30 kDa have been described. There is one splice variant that shows a 72 aa substitution for aa 1-106. Over aa 556-724, human SARM1 shares 94% aa identity with mouse SARM1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.