

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Caspase-8 precursor in Western blots and a 42 kDa doublet generated during apoptosis.
Source	Monoclonal Mouse IgG _{2A} Clone # 84131
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human Caspase-8 aa 234-391
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Caspase-8 (Cysteine-aspartic acid protease 8/Casp8; also MCH5 and FLICA) is a 28 kDa member of the peptidase C14A family of enzymes. It is known as the initiating caspase for the apoptotic cascade. Caspase-8 acts on procaspases-3, 4, 6, 7, 9 and 10, in addition to c-FLIPL and procaspase-8 itself. Human procaspase-8α4 is a 58 kDa (predicted), 496 amino acid (aa) protein and contains two N-terminal death domains plus a catalytic site that utilizes His334Gly335 plus Cys377. Following death domain-containing receptor activation, proteolysis generates a 28 kDa heterodimer. This includes a p18 subunit (aa 234-391; or 217-374 of the 8a/standard form) noncovalently linked to a p10 subunit (aa 402-496; or 385-479 of the 8a/standard form). Association with another p18/p10 heterodimer generates active caspase-8. There are two common procaspase-8 isoform variants. One 53 kDa isoform (8b) shows a deletion of aa 103-134, while a second 55 kDa (8a) isoform shows a deletion of aa 103-134 with a 15 aa insertion after Lys183. The p18 and p10 subunits are 68% and 82% aa identical, human to mouse, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.