

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse S1P ₁ /EDG-1 peptide in direct ELISAs.
Source	Monoclonal Rat IgG _{2A} Clone # 713412
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse S1P ₁ /EDG-1 synthetic peptide (T4-H28) Accession # O08530
Conjugate	Biotin Excitation Wavelength: N/A nm Emission Wavelength: N/A nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

S1P₁ (sphingosine 1-phosphate receptor-1), also known as EDG-1 (endothelial differentiation, G-protein coupled receptor-1) or S1PR1 (sphingosine-1-phosphate receptor 1) is a widely expressed, 37-40 kDa, G protein coupled receptor within the S1P subfamily of the EDG family. S1P₁ is one of five receptors for the bioactive lipid S1P and mediates most of S1P effects on angiogenesis, vascular maturation, and cell migration, especially T cell egress from lymphoid organs. Human and mouse S1P₁ share 84% amino acid identity within the N-terminal extracellular portion used as an immunogen.