

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CEBP α in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human CEBP beta, gamma, delta, epsilon, or zeta is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 692716
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human CEBP α Met1-Ala124 Accession # P49715
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CCAAT-enhancer binding protein alpha (CEBP α) is a widely expressed 42 kDa basic-leucine zipper transcription factor that forms functional homodimers and heterodimers with CEBP β and CEBP γ . CEBP α inhibits mitotic progression through multiple mechanisms and also promotes terminal differentiation of granulocytes and adipocytes. Its activity is regulated by SUMOylation and by phosphorylation at multiple serine residues. CEBP α contains three TE transactivation domains (aa 1-69, aa 70-96, and aa 126-200) and one basic DNA binding motif and leucine zipper (aa 286 - 345). A 30 kDa isoform is generated by the use of an internal translation initiation site. The p30 isoform functions as a dominant negative factor by heterodimerizing with and inhibiting full length CEBP α activity. Within aa 1-124, human CEBP α shares 93% amino acid sequence identity with mouse and rat CEBP α , respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.