

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human Pannexin-1 in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant human Pannexin-2 or -3 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 720505
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Pannexin-1 Pro298-Asp376 Accession # Q96RD7
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Pannexin-1 (PANX1) is an approximately 45 kDa member of the Pannexin family of four-transmembrane channel proteins with a conserved pattern of cysteines. It is expressed on epithelial cells, neuronal and glial cells, cardiac myocytes, T cells, and erythrocytes. Pannexin-1 forms large pore hexameric channels or heteromeric channels with Pannexin-2 and enables the efflux of ATP and UTP. Its conductance is increased in response to hypoxia, isotonic stress, NMDA R activation, and Caspase-3 and -7 activation during apoptosis. Extracellular release of ATP through Pannexin-1 activates P2X7 receptors which in turn induces the closure of Pannexin-1 channels. Pannexin-1 cooperates with P2X7 for inflammasome activation, Caspase-1 activation, and the release of mature IL-1β and also regulates T cell activation at the immunological synapse. Within aa 298-376 (in the C-terminal cytoplasmic domain), human Pannexin-1 shares 96% and 98% aa sequence identity with mouse and rat Pannexin-1, respectively. Alternate splicing generates an additional isoform that lacks the C-terminal four amino acids in the cytoplasmic tail.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.