

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human SHC1 in direct ELISAs and human, mouse, and rat SHC1 in Western blots. In direct ELISAs, no cross-reactivity with recombinant human SHC3 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 725001
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human SHC1 Pro379-Val470, Trp488-Val579 Accession # P29353
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SH2 domain-containing transforming protein C1 (SHC1), also known as ShcA, is a cytoplasmic adaptor protein that is important in the signal transduction from growth factor, cytokine, and lymphocyte antigen receptors. SHC1 contains a PTB/PID domain (aa 156-339), a collagen homology domain (aa 340-487), and an SH2 domain (aa 488-579). Alternate splicing generates additional isoforms that differ in the extent of N-terminal truncation. Activation of SHC1 by phosphorylation at Ser239, Ser240, and Tyr317 enables Shc1 to interact with the GRB2/SOS complex, leading to the transcription of genes involved in mitogenesis and apoptosis. The p46 and p52 isoforms promote mitogenic signaling, while the p66 isoform promotes apoptosis and functions as a negative regulator of SHC1-mediated mitogenic signal transduction. Within aa 488-579, human SHC1 shares 100% aa sequence identity with mouse and rat SHC1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.