

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Metadherin in direct ELISAs.
Source	Monoclonal Rat IgG _{2B} Clone # 754523
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse Metadherin Lys167-Ser297 Accession # Q80WJ7
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Metadherin (MTDH; also Lyric and Astrocyte-elevated gene-1/AEG1) is a unique molecule originally discovered to be upregulated in astrocytes following HIV infection. Although its predicted MW is 64 kDa, it runs anomalously at approximately 80 kDa in SDS-Page. It is widely expressed, and appears to be a component of the ER, nucleolus and inner nuclear membrane. Metadherin is reported to promote Akt/PI3 kinase activity, and suppress apoptosis-associated FOXO3A transcription. Mouse Metadherin is a type III (i.e.- a type I with no signal sequence) transmembrane protein 579 amino acids (aa) in length. It contains a luminal N-terminus (aa 1-48) with a 510 aa cytoplasmic C-terminus. There are no identifiable structural motifs, although two poly-Lys segments and two utilized Ser phosphorylation sites exist. Multiple bands at 80 kDa, 75 kDa, 50-55 kDa and 37 kDa are seen for Metadherin on SDS-Page. They may reflect the presence of at least two potential isoform variants that show 1) a deletion of aa 422-457, and 2) an Ile substitution for aa 190-579. Over aa 167-297, mouse Metadherin shares 98% and 97% aa sequence identity with rat and human Metadherin, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.