

## Human CLEC-2A Alexa Fluor® 405-conjugated Antibody

Monoclonal Mouse IgG<sub>1</sub> Clone # 705518

Catalog Number: FAB7219V

DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human CLEC-2A in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) CLEC-2D, rhCLEC-2, or rhCD69 is observed.		
Source	Monoclonal Mouse IgG <sub>1</sub> Clone # 705518		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CLEC-2A Trp49-Leu174 (predicted) Accession # Q6UVW9		
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm		
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.		

Αŀ	'PL	ICA	IIOI	45

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 <sup>6</sup> cells	U937 human histiocytic lymphoma cell line

## PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze.

12 months from date of receipt, 2 to 8 °C as supplied.

## BACKGROUND

CLEC-2A (C-type lectin domain family 2 member A), also known as PILAR and KACL, is an approximately 35 kDa type II transmembrane glycoprotein in the C-type lectin superfamily. It contains one C-type lectin domain in its extracellular region. Alternate splicing of human CLEC-2A generates an additional isoform with a substitution of 37 amino acids (aa) at the C-terminus. CLEC-2A is upregulated on antigen-activated CD8<sup>+</sup> T cells and enhances T cell proliferation through interactions with CD161. It is also expressed on keratinocytes and binds to NKp65/KLRF2, triggering NK cell activation.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

