

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Attractin in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 750003
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Attractin Ala84-Gln1272 Accession # O75882
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Attractin (ATRN), also known as DPPT-L, is an approximately 200 kDa transmembrane glycoprotein that shows dipeptidyl peptidase activity similar to DPPIV/CD26. Attractin is involved in a variety of processes including monocyte-T cell adhesion, axon myelination, melanocyte pigment synthesis, and energy homeostasis. The extracellular region of human Attractin contains one EGF-like domain, one CUB domain, six Kelch repeats, four PSI domains, one C-type lectin domain, and two laminin EGF-like domains. Alternate splicing of human Attractin generates a secreted isoform that lacks the transmembrane and cytoplasmic regions. Attractin is transiently upregulated during T cell activation before expression switches to the 175 kDa secreted isoform which is released into the circulation. Soluble Attractin is preferentially expressed by leukocytes and differentiating neurons. It blocks neurite formation and is elevated in the CSF of astrocytoma patients. Within aa 84-1272, human Attractin shares 95% aa sequence identity with mouse and rat Attractin.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.