

## DESCRIPTION


|                           |  |
|---------------------------|--|
| <b>Species Reactivity</b> | Mouse  |
| <b>Specificity</b>        | Detects mouse CD79B in direct ELISAs.  |
| <b>Source</b>             | Monoclonal Rat IgG <sub>2A</sub> Clone # 735451  |
| <b>Purification</b>       | Protein A or G purified from hybridoma culture supernatant   |
| <b>Immunogen</b>          | Chinese hamster ovary cell line CHO-derived recombinant mouse CD79B<br>Met29-Asp158<br>Accession # P15530  |
| <b>Conjugate</b>          | Phycoerythrin<br>Excitation Wavelength: 488 nm<br>Emission Wavelength: 565-605 nm  |
| <b>Formulation</b>        | Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.<br><br>*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

|                       | Recommended Concentration        | Sample    |
|-----------------------|----------------------------------|-----------|
| <b>Flow Cytometry</b> | 10 $\mu$ L/10 <sup>6</sup> cells | See Below |

## DATA



## PREPARATION AND STORAGE

|                                |  |
|--------------------------------|--|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.                                  |
| <b>Stability &amp; Storage</b> | <b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul> |

## BACKGROUND

CD79B (also known as B29, Ig $\beta$  and B cell antigen receptor complex-associated protein  $\beta$ -chain) is a 37-39 kDa member of the Ig-Superfamily. It is expressed on B cells and forms a covalent heterodimer with CD79A. This complex interacts noncovalently with membrane Ig, forming the B cell antigen receptor. Within this complex, membrane Ig detects antigen while CD79A:B initiates signaling. CD79B is also required for formation of pre-B cells during B cell development. Mature mouse CD79B is a 203 amino acid (aa) type I transmembrane glycoprotein (aa 26-228). It contains an extracellular region with one V-type Ig-like domain (aa 41-132) and an ITAM-containing cytoplasmic domain (aa 181-228). CD79B may migrate as two bands in SDS-PAGE. One defines the standard 37 kDa form, while a second represents one of two possible isoforms, the first of which is an underglycosylated full-length CD79B, and the second of which is an alternative splice form that likely lacks the C-terminal 32 amino acids. Mouse CD79A and CD79B share only 24% aa identity. Over aa 29-158, mouse CD79B shares 54% and 78% aa identity with human and rat CD79B, respectively.