

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CDK5 Activator 1 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human CDK5 Activator 2 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 739304
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human CDK5 Activator 1 Ser8-Phe98 Accession # Q15078
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cyclin dependent kinase 5 regulatory subunit 1 (CDK5R1) is a 35 kDa protein that associates with and activates CDK5. Its expression is restricted to neurons and is required for normal brain development. CDK5R1 can be cleaved by calpain at Phe98-Ala99 which liberates a C-terminal fragment known as p25. p25 induces prolonged activation of CDK5 and broadens the substrate specificity of CDK5 to include tau which is aberrantly phosphorylated in Alzheimer's disease. Full length CDK5R1 is myristoylated at Gly2 and is preferentially localized to membranes at the cell periphery. p25 lacks this modification and is enriched in the nucleus and perinuclear regions. p25 accumulates in the brains of Alzheimer's patients where it promotes neuronal apoptosis and cytoskeletal abnormalities. Within aa 100-307, human CDK5R1 shares 99% aa sequence identity with mouse and rat CDK5R1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.