

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse VEGF R3/Flt-4 in direct ELISAs and Western blots. In direct ELISAs, approximately 20% cross-reactivity with recombinant human (rh) VEGF R3/Flt-4, 10% cross-reactivity with rhVEGF R2, and 5% cross-reactivity with rhVEGF R1.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant mouse VEGF R3/Flt-4 Tyr25-Asp770 Accession # P35917
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

VEGF R3 (Flt-4), together with VEGF R1 (Flt-1) and VEGF R2 (KDR/Flk-1), belong to the class III subfamily of receptor tyrosine kinases (RTKs). All three receptors contain seven immunoglobulin-like repeats in their extracellular domains and kinase insert domains in their intracellular regions. The expression of these receptors is almost exclusively restricted to the endothelial cells. These receptors are likely to play essential roles in vasculogenesis and angiogenesis.

In adults, VEGF R3 expression is restricted to the endothelial cells of the lymphatic vessels. Mouse VEGF R3 cDNA encodes a 1363 amino acid (aa) residue precursor protein with a 24 aa residue signal peptide. Mature VEGF R3 has a 751 aa residue extracellular domain, a 22 aa residue hydrophobic transmembrane domain and a 565 aa residue cytoplasmic domain. The polypeptide sequences of murine Flt-4 is 88% identical to the human homologue. VEGF R3 has been reported to serve as the receptors for VEGF-C and VEGF-D.

References:

1. Finnerty, H. *et al.* (1993) *Oncogene* **8**:2293.
2. Joukov, V. *et al.* (1996) *EMBO J.* **15**:290.
3. Achen, M. *et al.* (1998) *Proc. Natl. Acad. Sci. USA* **95**:548.