

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human CD1b in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) CD1a, rhCD1c, rhCD1d, rhCD1e, recombinant mouse (rm) CD1d1, or rmCD1d2 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 737249
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CD1b Ser18-Ser303 Accession # P29016
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	5 µL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CD1b is a 44-46 kDa member of the CD1 family of proteins. It is one of five CD1 isoforms (CD1a-e) that resemble MHC-I type molecules. Based on structure, CD1a, 1b and 1c can be grouped together, with CD1d and 1e existing as stand-alone forms. CD1b is a 316 amino acid (aa) type I transmembrane glycoprotein that contains one Ig-like domain in its ECD. It is expressed in the late endosome/lysosome compartment of myeloid-type dendritic cells. Following synthesis, CD1b associates with SAP-C, binds two short lipids in an unusually large ligand-binding groove, and transits either to the plasma membrane, or more commonly, the endosomal compartment. In either case, exposure to Gm⁻ lipid A or mycobacterial glycolipid results in a substitution of the foreign lipid for the two short-chain self lipids, followed by its presentation on the plasma membrane. Select αβ and γδ T cell recognize these lipids and typically respond with Th1-type cytokines (IFN-γ and TNF-α). Unlike other CD1 family molecules, CD1b does not seem to have a soluble form. There is no rodent counterpart to human CD1b, only to CD1d.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.