

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CD161 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 702228
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	BaF3 mouse pro-B cell line transfected with human CD161 Accession # Q12918
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human peripheral blood lymphocytes

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CD161, also known as NKR-P1A and KLRB1, is a 40 kDa type II transmembrane glycoprotein that contains one C-type lectin domain in its extracellular region. CD161 is expressed as a disulfide-linked dimer on the surface of Th17 cells and NK cells as well as on subsets of CD1-restricted T cells, intestinal NT cells, peripheral memory T cells, monocytes, and dendritic cells. It binds to OCIL/CLEC2d, leading to an inhibition of NK cell-mediated cytotoxicity and IFN-γ secretion. Alternatively, CD161 can enhance TCR activation to CD1d ligation. CD161⁺ cell populations are depleted in ulcerative colitis, Grave's disease, and AIDS, although CD161⁺ T cells are activated during asthmatic attacks. Additional related proteins are expressed in mouse but not human: the inhibitory NKR-P1B and D, and the stimulatory NKR-P1A, C, and F. Human CD161 shares 47% aa sequence identity with mouse and rat NKR-P1A.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.