

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Carboxypeptidase M in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG _{2B} Clone # 848543
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Carboxypeptidase M Leu18-Ser423 Accession # P14384
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Carboxypeptidase M (CPM) is a 50-65 kDa monomer that belongs to the regulatory CPN/E subfamily, peptidase M14 family of enzymes. It is widely expressed, being found on macrophages, fibroblasts, endothelial cells, oligodendrocytes and Schwann cells, dendritic cells, osteoblasts and bronchial epithelium. Carboxypeptidase M is a GPI-linked glycoprotein that is best known as a peptidase that cleaves basic amino acids (aa) from the carboxyterminal of a number of peptides, including EGF and bradykinin. It is also known to bind to apparent substrates and undergo a conformational change that links it with the kinin B1 GPCR, initiating signal transduction. Mature human Carboxypeptidase M is 406 aa in length (aa 18-423). It contains one large enzymatic region (aa 19-310) and two critical glutamic acid residues at Glu260 and Glu264. Like many GPI-linked proteins, Carboxypeptidase M undergoes solubilization and is reportedly found in urine and amniotic fluid. Over aa 18-423 (mature Carboxypeptidase M), human Carboxypeptidase M shares 86% aa sequence identity with mouse Carboxypeptidase M.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.