

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Peroxiredoxin 3 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human Peroxiredoxin 1, 2, 4, or recombinant mouse Peroxiredoxin 3 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 774406
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Peroxiredoxin 3 Met1-Gln256 Accession # P30048
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Peroxiredoxin 3 (Prx-3/III; also AOP-1, MER5 and thioredoxin-dependent peroxidase reductase) is a ubiquitous, 22-28 kDa mitochondrial antioxidant enzyme that belongs to the 2-Cys class of the TSA/ahpC family of peroxiredoxins. Prx-3 is known to act as either a homodimer, or a decamer, and scavenge reactive oxygen species generated by oxidative stress. The mouse Prx-3 precursor molecule is 257 amino acids (aa) in length. It contains a cleavable N-terminal 63 aa mitochondrial targeting sequence, plus a 194 aa mature enzyme that shows a thioredoxin domain between aa 64-222. There are two catalytic cysteines, one at Cys109 and another at Cys230 of the precursor. Prx-3 undergoes a phosphorylation at Thr147 that reduces its activity. One potential splice form shows a deletion of Arg149, Lys150 and Arg185. Full-length mouse Prx-3 is 86% and 95% aa identical to human and rat Prx-3, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.