

Mouse Endosialin/CD248 Alexa Fluor® 488-conjugated Antibody

Monoclonal Rat IgG_{2A} Clone # 458606 Catalog Number: FAB7535G

100 µg

DESCRIPTION	
Species Reactivity	Mouse
Specificity	Detects mouse Endosialin/CD248 in direct ELISAs. In direct ELISAs, less than 10% cross-reactivity with recombinant human Endosialin/CD248 is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 458606
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Endosialin/CD248 Val13-Arg693 (predicted) Accession # Q91V98
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

China | info.cn@bio-techne.com TEL: 400.821.3475

PREPARATION AND STORAGE	
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Endosialin, also known as CD248 and tumor endothelial marker 1 (Tem1), is a 165 kDa transmembrane O-glycosylated protein that contains one C-type lectin, one Sushi, one EGF-like domain and a mucin-like stalk in its extracellular domain (ECD). It is expressed on activated perivascular and stromal cells in embyronic and tumor neovasculature but is downregulated in quiescent vasculature. Endosialin regulates pericyte proliferation, migration, and adhesion to matrix Fibronectin and Collagens I and IV. It is also required for lymph node and spleen remodeling during immune responses. Within the ECD, mouse Endosialin shares 76% and 93% aa sequence identity with human and rat Endosialin, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/23/2025 Page 1 of 1