

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human BMP-4 in direct ELISAs and Western blots. This antibody shows approximately 5% cross-reactivity with recombinant human BMP-2.
Source	Monoclonal Mouse IgG _{2B} Clone # 66119
Purification	Protein A or G purified from ascites
Immunogen	Mouse myeloma cell line NS0-derived recombinant human BMP-4 Ser293-Arg408 Accession # P12644
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

BMP4 is a TGFβ superfamily ligand that is widely expressed from early embryogenesis through adulthood. It plays an important role in mesenchyme formation, epidermal determination, suppression of neural induction, the development of multiple organs, and tissue repair (1-5). The human BMP4 precursor contains a 273 amino acid (aa) propeptide and a 116 aa mature protein (6). Processing of the propeptide by furin or proprotein convertase 6 enables the formation of the mature disulfidelinked homodimeric BMP4 and facilitates its secretion. Similar intracellular processes may lead to the formation and recreation of BMP4/BMP7 disulfidelinked heterodimer (7-9). Mature human and mouse BMP4 share 98% aa sequence identity. Human BMP4 shares 85% aa sequence identity with human BMP2 and less than 50% with other human BMPs. Compared to BMP4 homodimers, BMP4/ BMP7 heterodimers exhibit a greater potency in inducing osteogenic differentiation (9). In *Xenopus*, the heterodimers can also induce the formation of mesoderm, whereas BMP4 homodimers only provide ventralizing signals for existing mesoderm (10). BMP4 signals through tetrameric complexes composed of type I (primarily Activin RIA or BMPRIA) and type II (primarily Activin RIIA or BMPRII) receptors (11, 12). The bioavailability of BMP4 is regulated by its interaction with multiple proteins and glycosaminoglycan (13-15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.