

#### DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human NAALADase-2/NAALAD2 in direct ELISAs. In direct ELISAs, approximately 50% cross-reactivity with recombinant mouse NAALADase-2 is observed and no cross-reactivity with recombinant human (rh) PSMA/FOLH1/NAALADase-1, rhNAALADase-like 2, or rh
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 756030
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human NAALADase-2/NAALAD2 Lys32-Leu740 Accession # Q9Y3Q0
<b>Conjugate</b>	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
<b>Formulation</b>	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

<b>Western Blot</b>	Optimal dilution of this antibody should be experimentally determined.
<b>Immunohistochemistry</b>	Optimal dilution of this antibody should be experimentally determined.

#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

#### BACKGROUND

N-acetylated-alpha-linked acidic dipeptidase 2 (NAALADase-2), also known as glutamate carboxypeptidase 3 (GCP3), is a 90-100 kDa type 2 transmembrane protein in the M28 peptidase family. It hydrolyzes beta-citrylglutamate (BCG) and N-acetyl-aspartylglutamate (NAAG) which are abundant dipeptides found in brain and testis. Mature human NAALADase-2 consists of a 7 amino acid (aa) cytoplasmic domain, a 24 aa transmembrane segment, and a 709 aa extracellular region (aa 32-740) that contains the peptidase domain. Over aa 32-740, human NAALADase-2 shares 89% aa sequence identity with mouse NAALADase-2.

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.