

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse TIMP-4 in direct ELISAs.
Source	Monoclonal Rat IgG ₁ Clone # 866106R
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse TIMP-4 Met1-Pro224 Accession # Q9JHB3
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Tissue Inhibitors of Metalloproteinases (TIMPs) are a family of secreted proteins that regulate the activation and proteolytic activity of the zinc enzymes known as Matrix Metalloproteinases (MMPs). There are four known members of the family, TIMP-1, -2, -3, and -4. TIMP-4 is produced by a wide range of tissues, particularly brain, heart, ovary and skeletal muscle (1, 2). Limited studies have shown that TIMP-4 has a tumor suppressive effect against Wilm's tumor, exhibits negative correlation with glioma malignancy and is found in breast carcinoma cells (3, 5). TIMP-4 inhibits MMP-mediated proteolysis by forming a noncovalent binary complex with the MMP active site through its N-terminal domain. In addition, it binds to the hemopexin-like domain of proMMP-2 through its C-terminal domain in a manner similar to TIMP-2 (6). However, unlike TIMP-2, TIMP-4 does not promote proMMP-2 activation by MT1-MMP (MMP-14) (7). Although TIMP-4 is a potent inhibitor of most MMPs, it is not an effective inhibitor of ADAMs, such as TACE (8, 9).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.