

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human BetaTrCP1 isoform 2 in direct ELISAs and Western blots. Detects mouse BetaTrCP1 in Western Blots. Detection of mouse BetaTrCP1 in Immunocytochemistry has not been tested.
Source	Monoclonal Mouse IgG ₁ Clone # 763524
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human β -TrCP1 isoform 2 Met1-Leu120 Accession # Q9Y297-2
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

β -TrCP1, also called BTRC (β -transducin repeat containing E3 ubiquitin ligase), FBW1A (F-box/WD repeat containing 1A) and FBXW1A, is an approximately 63-70 kDa protein that associates with Skp1, Cullin, and RBX1 to form an SCF ubiquitin ligase complex. These complexes regulate the degradation of a wide variety of proteins including transcription factors, signal transduction molecules, and cytokine receptors. They play important roles in regulation of the inflammatory response, cell cycle, apoptosis, tumorigenesis, and the response to DNA damage. β -TrCP1 can form homodimers and heterodimers with β -TrCP2, which associate into SCF complexes with distinct substrate preferences. Human β -TrCP1 contains a dimerization domain (aa 128-177), an F-box domain (aa 181-228), and seven tandem WD repeats (aa 301-590). The WD repeats assemble into a beta propeller structure that binds to phosphorylated destruction motifs in the substrate. Alternate splicing generates a short isoform with a 36 aa deletion near the N-terminus. Within aa 1-120 of the 63 kDa short isoform, human β -TrCP1 shares 98% and 91% aa sequence identity with mouse and rat β -TrCP1, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.