

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human ALK-7 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant rat ALK-7 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 810506
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human ALK-7 Leu26-Glu113 Accession # Q8NER5
<b>Conjugate</b>	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	10 $\mu$ L/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

Activin Receptor-like Kinase 7 (ALK-7), also known as Activin R1C (gene name ACVR1C), is a glycosylated 58 kDa type I receptor in the superfamily of TGF- $\beta$  serine/threonine kinase receptors. It associates with type II receptors to form a signaling complex that responds to the ligands Activin AB, and Activin B, GDF3, and Nodal. ALK-7 plays a role in regulating energy balance by inhibiting insulin secretion and inducing pancreatic beta cell apoptosis. It is expressed in adipose tissue but downregulated in obesity. ALK-7 is also expressed in pituitary gonadotrophic cells and in pre-eclamptic placenta. It induces the apoptosis of trophoblasts as well as ovarian granulosa and epithelial cells. Within the extracellular domain, human ALK-7 shares 95% and 91% amino acid (aa) sequence identity with mouse and rat ALK-7, respectively. Alternate splicing of human ALK-7 generates additional isoforms with either a 50 aa N-terminal truncation or with deletions of 79 aa or 157 aa that encompass the transmembrane segment.