

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Lgr4/GPR48 in Lgr4-transfected cell-based ELISA. Stains human Lgr4/GPR48 transfected but not irrelevant transfected cells in flow cytometry.
Source	Monoclonal Mouse IgG _{2B} Clone # 852229
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with human Lgr4/GPR48 Met1-Asp951 Accession # Q9BXB1
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	HEK293 human embryonic kidney cell line transfected with human Lgr4/GPR48 and eGFP

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Lgr4 (Leucine-rich repeat GPR 4), also known as GPR48 (G-Protein-coupled Receptor 48), is a seven-transmembrane glycoprotein receptor in the Lgr family of cell surface receptors (1, 2). While this family includes receptors for hormones such as LH, FSH, TSH, and HCG, the subfamily comprising Lgr4, Lgr5, and Lgr6 are G-protein-independent mediators of the potentiating effect of R-Spondins on Wnt signaling (1-6). Lgr4 binds and forms complexes with R-Spondins, Frizzled Wnt receptors and LRP Wnt co-receptors (5). It acts at least in part by enhancing Wnt-dependent LRP phosphorylation, internalization of LRPs, and accumulation of β-catenin (3, 4). Human Lgr4 cDNA encodes 951 amino acids (aa), including a long N-terminal Extracellular Domain (ECD, aa 25-544) with 16-17 LRR domains that mediate ligand interaction (1). The LRR-containing ECD of human Lgr4 shares 93% aa sequence identity with mouse, rat and bovine Lgr4, and 50-60% aa identity with human Lgr5 and Lgr6. Lgr4 is widely expressed in both embryo and adult. Expression of Lgr4 mRNA in adult humans is highest in pancreas, followed by liver, heart, muscle, brain, and placenta (1). In rodents, embryonic and adult expression includes liver, kidney, adrenals, bone/cartilage, and heart (2, 7-9). Lgr4 deletion in the mouse affects development in areas of expression, for example, inhibiting fetal liver definitive erythropoiesis (9). Deletion of Lgr4 specifically from stem and progenitor cells in intestinal crypts induces loss of crypts due to insufficient Wnt signaling (5, 6). Lgr4 may be over-expressed in carcinomas and may promote invasiveness and metastasis by down-regulating p27^{Kip1} expression (10).

References:

1. Loh, E.D. *et al.* (2001) *Biochem. Biophys. Res. Commun.* **282**:757.
2. Hsu, S.Y. *et al.* (1998) *Mol. Endocrinol.* **12**:1830.
3. Carmon, K.S. *et al.* (2011) *Proc. Natl. Acad. Sci. USA* **108**:11452.
4. Glinka, A. *et al.* (2011) *EMBO Rep.* **12**:1055.
5. de Lau, W. *et al.* (2011) *Nature* **476**:293.
6. Ruffner, H. *et al.* (2012) *PLoS ONE* **7**:e40975.
7. Van Schoore, G. *et al.* (2005) *Histochem. Cell Biol.* **124**:35.
8. Mazerbourg, S. *et al.* (2004) *Mol. Endocrinol.* **18**:2241.
9. Song, H. *et al.* (2008) *J. Biol. Chem.* **283**:36687.
10. Gao, Y. *et al.* (2006) *Cancer Res.* **66**:11623.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Human Lgr4/GPR48
Alexa Fluor[®] 700-conjugated Antibody

Monoclonal Mouse IgG_{2B} Clone # 852229

Catalog Number: FAB7750N
100 µg