

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human L1CAM in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) VCAM-1, rhICAM-1, rhICAM-2, and rhICAM-3 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 84321
Purification	Protein A or G purified from ascites
Immunogen	Mouse myeloma cell line NS0-derived recombinant human L1CAM Met1-Glu1120 Accession # CAA42508
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

L1CAM (Neural cell adhesion molecule L1, also known as L1, CD171 and NCAM-L1) is a 200-230 kDa member of the L1 family, Immunoglobulin (Ig) superfamily of molecules. L1 is recognized to play a key role in cell migration, adhesion, neurite outgrowth, myelination and neuronal differentiation. It does so through a series of *cis* and *trans* interactions that involve multiple copartners and target receptors. L1 is described as forming both homotypic and heterotypic complexes, the latter with molecules as diverse as the EGFR, NCAM, CD24, neurocan and various α _v plus β ₁ and β ₃ integrins. Cells known to express L1 include immature oligodendrocytes, CD4+ T cells, B cells and monocytes, premyelinating Schwann cells, intestinal epithelial progenitor cells, and cerebellar granule plus Purkinje cells. Mature human L1 is a 1238 amino acid (aa) type I transmembrane protein. It contains an 1101 aa extracellular region (aa 20-1120) plus a 114 aa cytoplasmic domain (aa 1144-1257). The extracellular region possesses six C2-type Ig-like domains (aa 35-607) followed by five fibronectin (FN) type III repeats (aa 612-1108). The cytoplasmic tail contains no kinase motifs, but does possess a FIGQY peptide that interacts with ankyrin, and an RSLE sequence that mediates clathrin-associated endocytosis. At least five Ser residues are known to be phosphorylated. There are two splice variants, one each in the intracellular and extracellular domain. A deletion of RSLE (aa 1177-1180) adversely affects endocytosis, while a Leu substitution for aa 26-31 interferes with numerous heterotypic interactions. In general, the full-length L1 molecule is a neuron-associated isoform. L1 is known to undergo proteolysis, either by plasmin or ADAMs. This generates soluble isoforms of varying sizes (140-200 kDa) that retain bioactivity, and which can be incorporated into the surrounding ECM. The membrane fragments (30-80 kDa) undergo further processing, most importantly by γ -secretase, to generate a soluble 28 kDa intracellular domain. This domain is SUMOylated, and believed to possess an NLS at Lys1147. Upon presumed entry into the nucleus, L1 is posited to activate L1-responsive genes. In the extracellular region, human and mouse L1 share 86% aa sequence identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.