

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Protocadherin-12 in ELISA.
Source	Monoclonal Rat IgG _{2A} Clone # 868909
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Protocadherin-12 Met1-Ala716 Accession # O55134
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

PCDH12 (Protocadherin 12; also VE-Cadherin 2) is a 150-160 kDa glycoprotein member of the PCDH family of molecules. It exhibits limited expression, being found on endothelial cells, renal mesangium, Sertoli cells, and glycogen-positive trophoblast cells. PCDH is a transmembrane protein that forms homotypic aggregates in a calcium-dependent manner. Although this would suggest a prominent role in cell-cell adhesion, it is unclear if this is the only activity for the molecule. In any case, it appears that PCDH12 does play an important role in placental development. Mature mouse PCDH12 is an 1163 amino acid (aa) type I transmembrane protein (aa 18-1180). It possesses a 699 aa extracellular region (aa 18-716) that contains six cadherin domains (aa 28-711), plus a 21 aa transmembrane segment coupled to a lengthy 443 aa cytoplasmic domain (aa 738-1180). Based on human, mouse PCDH12 will undergo proteolytic processing. MMP activity will first generate a 90 kDa circulating extracellular fragment and a 60 kDa membrane-bound fragment, followed by γ-secretase which has the potential to further act on the membrane-bound fragment, generating a cytosolic 50 kDa fragment. There is one potential alternative start site at Met1060. Over aa 18-716, mouse PCDH12 shares 95% and 82% aa sequence identity with rat and human PCDH12, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.