

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Casein Kinase 2α in ELISAs and Western blots. Detects mouse and rat Casein Kinase 2α in Western blots
Source	Monoclonal Mouse IgG ₁ Clone # 844720
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Casein Kinase 2α Asp253-Gln391 Accession # P68400
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Knockout Validated	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Casein kinase 2 (CK2) is a ubiquitous and constitutively active tetrameric serine/threonine kinase that is comprised of two catalytic subunits (CK2α and/or CK2α') and two identical regulatory subunits (CK2β). CK2 has been implicated in numerous cellular processes, including signal transduction, transcription, translation, replication, and metabolic pathways. CK2 is known to phosphorylate more than 300 different substrates. Phosphorylation of cell-cycle proteins such as p53, p34cdc2, p27KIP1, and MDM-2 account for the ability of CK2 to induce proliferation, while the phosphorylation of HS1, Bid, and Max account for its antiapoptotic role. The human CK2α and CK2α' subunits are the products of two different genes. They have highly conserved catalytic domains but divergent C-terminal regions. Within aa 253-391 (including the region of divergence between CK2α and CK2α'), the 35-45 kDa human CK2α shares 96% aa sequence identity with mouse and rat CK2α. An alternatively spliced isoform of human CK2α lacks the N-terminal 136 amino acids including a portion of the kinase domain. CK2β plays dual roles in the regulation of CK2 activity. Its C-terminal domain is responsible for stable interactions with the catalytic subunit and increased catalytic activity following tetramer formation, while the N-terminal domain exerts negative regulation on the catalytic activity of CK2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.