

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Nephronectin in ELISAs and Western blots.
Source	Monoclonal Mouse IgG _{2A} Clone # 859326
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Nephronectin Glu20-Arg565 Accession # Q6UXI9
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Nephronectin, also called POEM (preosteoblast EGF repeat protein with MAM domain) or EGFL6L, is a 70-90 kDa extracellular matrix protein that is a ligand for integrin α8β1 and mediates cell adhesion and spreading. The 565 amino acid (aa) human Nephronectin contains a 19 aa signal sequence, five EGF-like domains, a mucin-like region, an RGD integrin binding motif, and a MAM domain. Isoforms of 509-595 aa contain deletions of aa 387-416, and/or an insertion of 30 aa after aa 88, and/or substitution of 3 aa for aa 535-564. Mature human Nephronectin shares 88% and 89% aa sequence identity with mouse and rat Nephronectin, respectively. It is most highly expressed in developing endocrine organs such as parathyroid, thyroid, hypophysis and pineal organ, around developing bone, teeth, and muscle, and in the Wolffian duct and ureteric bud basement membranes in the developing kidney.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.