

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human Kynurenine 3-Monooxygenase/KMO in direct ELISAs. Detects human, mouse, and rat Kynurenine 3-Monooxygenase/KMO in Western blot.
Source	Monoclonal Rabbit IgG Clone # 2493A
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human Kynurenine 3-Monooxygenase/KMO Asp2-Leu441 Accession # O15229
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Kynurenine 3-Monooxygenase (KMO), also known as Kynurenine 3-Hydroxylase, is a part of the kynurenine pathway of tryptophan degradation (1). KMO catalyzes the NADPH- and flavin adenine dinucleotide (FAD)-dependent 3-hydroxylation of kynurenine to 3-hydroxykynurenine (3-HK). 3-HK is neurotoxic via the generation of hydrogen peroxide (2) and through the excitotoxic effects of its downstream metabolite quinolinic acid (3). The levels of 3-HK and quinolinic acid are increased in the brain with Alzheimer's disease and Huntington's disease (1). Inhibition of KMO was shown to reverse cognitive and motor deficits in mouse models of those diseases via an increase in neuroprotective kynurenic acid (4). KMO is found in the mitochondrial outer membrane of microglial cells in the brain and dendritic cells and macrophages in the periphery (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.