

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human DAB2 in ELISA and Western Blot.
Source	Monoclonal Mouse IgG ₃ Clone # 883216
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human DAB2 Lys630-Ala770 Accession # P98082
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

DAB2 (Drosophila disabled homolog 2; also DOC-2/Differentially-expressed in Ovarian Carcinoma and p96) is a cytoplasmic phosphoprotein. Although the predicted MW is 82 kDa, it runs anomalously at 95-105 kDa in SDS-PAGE. It is widely expressed, being found in ovarian cuboidal epithelium, megakaryocytes, fibroblasts, macrophages, breast epithelium and renal proximal tubule cells. Reduced expression levels are associated with tumor development. DAB2 has multiple functions, but is typically described as an adaptor protein (i.e. one that supports the apposition of two interacting molecules) involved in intracellular trafficking. Molecules reported to interact with DAB2 include megalin, SMAD2 plus TGF-βRI/II, axin and Dvl3. With respect to trafficking, DAB2 reportedly binds to AP-2, clathrin, LRP6 and myosin VI, four molecules associated with the endocytic process. Human DAB2 is 770 amino acids (aa) in length. It contains one pleckstrin homology-like domain that binds phosphotyrosine (aa 42-170) and a C-terminal Pro-rich region that binds SH3 domains (aa 600-630). There are two isoform variants. One is 80-82 kDa in size and shows a deletion of aa 230-447. The second exhibits a short deletion of aa 209-229. Over aa 630-770, human DAB2 shares 76% aa sequence identity with mouse DAB2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.