

DESCRIPTION

| | |
|---------------------------|--|
| Species Reactivity | Human |
| Specificity | Detects human B7-H7/HHLA2 in direct ELISAs. |
| Source | Monoclonal Mouse IgG ₁ Clone # 907812 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | Human embryonic kidney cell line HEK293-derived human B7-H7/HHLA2 Met1-Asn344 Accession # Q9UM44 |
| Conjugate | Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm |
| Formulation | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|----------------------------------|---|
| Flow Cytometry | 0.25-1 µg/10 ⁶ cells | HEK Human Cell Line Transfected with Human B7-H7/HHLA2 and eGFP |

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

B7-H7, also known as HHLA2 (HERV-H LTR-associating 2), is a member of the B7 family of immune regulatory proteins (1, 2). Mature human B7-H7 consists of a 322 amino acid (aa) extracellular domain (ECD) with three immunoglobulin-like domains, a 21 aa transmembrane segment, and a 49 aa cytoplasmic domain (3-5). B7-H7 is constitutively expressed on monocytes and is up-regulated by LPS and IFN-γ stimulation. It is expressed on LPS/IFN-γ treated B cells but not on resting B cells (5). B7-H7 binds to cell surface determinants on resting and mature T cells, B cells, and monocytes as well as on immature and mature dendritic cells (5). Soluble B7-H7 inhibits the proliferation of activated CD4⁺ and CD8⁺ T cells and their production of IFN-γ, TNF-α, IL-5, IL-10, IL-13, IL-17A, and IL-22 (5).

References:

1. Zou, W. and L. Chen (2008) *Nat. Rev. Immunol.* **8**:467.
2. Bour-Jordan, H. *et al.* (2011) *Immunol. Rev.* **241**:180.
3. Mager, D.L. *et al.* (1999) *Genomics* **59**:255.
4. Flajnik, M.M. *et al.* (2012) *Immunogenetics* **64**:571.
5. Zhao, R. *et al.* (2013) *Proc. Natl. Acad. Sci. USA* **110**:9879.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.