

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Phospholipid Scramblase 1/PLSCR1 in ELISA.
Source	Monoclonal Mouse IgG ₁ Clone # 875327
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Phospholipid Scramblase 1/PLSCR1 Met1-Pro84 Accession # O15162
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

PLSCR1 (phospholipid scramblase 1) is an approximately 37 kDa, type II transmembrane protein that is a member of the Ca²⁺-dependent phospholipid scramblase family that regulates phospholipid asymmetry of membranes. It is most highly expressed on erythrocytes and platelets, and plays roles in exocytosis and apoptosis. PLSCR1 also has nuclear activity, for example, as an activator of genes in response to interferons. The 318 amino acid (aa) human PLSCR1 contains a 288 aa cytoplasmic domain with three SH3-binding and two WW-binding adaptor domains, three phosphorylation sites, four potential palmitoylation sites (regulating localization to the nucleus or plasma membrane) and a nuclear localization signal, followed by a transmembrane domain and a short extracellular domain. The N-terminus of human PLSCR1 (aa 1-84) shares <60% aa sequence identity with mouse and rat PLSCR1. Potential isoforms have a downstream start site without (311 aa) or with (186 aa) divergence after aa 192, or an upstream start site with divergence of 1 or 11 aa after aa 192 (205, 215 aa).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.