

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Hexokinase 2 in direct ELISAs. Detects human Hexokinase 1 and human Hexokinase 2 in Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 927312
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Hexokinase 2. Phe11-Arg917 Accession # P52789
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Hexokinases phosphorylate hexose to form hexose 6-phosphate, the first step in hexose metabolism (1). Phosphorylation of a hexose adds charge to molecule thereby making it difficult to transport out of a cell. The hexose is therefore retained for intracellular metabolic processes, such as glycolysis or glycogen synthesis. In most organisms, glucose is the most important substrate of hexokinases and glucose-6-phosphate is the most important product. There are four mammalian hexokinases (2). Hexokinase 1, 2 and 3 are referred to as high-affinity hexokinases because their Km for glucose is below 1 mM. Hexokinase 4 is specific for glucose and is also referred to as glucokinase (3). Hexokinase 2 (HK2), also known as muscle form hexokinase, localizes to the outer membrane of mitochondria and is present in adipose tissue, skeletal muscle, and heart (4). The amino acids corresponding to the mitochondrial binding domain (5) have been removed in the recombinant enzyme. Like Hexokinase 1 (HK1), HK2 contains two homologous halves that may have evolved from an ancestral hexokinase through gene duplication and tandem ligation (6). Unlike HK1, in HK2 both the C-terminal and N-terminal portions are catalytically active with the N-terminal half having higher activity than the C-terminal half (7). In HK2 both the N-terminal and C-terminal halves exhibit product inhibition. HK2 overexpression is required for tumor growth making HK2 an attractive oncotarget (4). The enzymatic activity of recombinant human HK2 is measured using a phosphatase-coupled method (8).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.