

## DESCRIPTION

|                           |                                                                                                                                                                                                                                                                             |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b> | Human                                                                                                                                                                                                                                                                       |
| <b>Specificity</b>        | Detects human XIAP in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) cIAP-1 or rh cIAP-2 is observed. Neutralizes human XIAP and human XIAP-BIR2 bioactivity. Does not neutralize XIAP-BIR3 or Livin bioactivity.                         |
| <b>Source</b>             | Monoclonal Mouse IgG <sub>1</sub> Clone # 117318                                                                                                                                                                                                                            |
| <b>Purification</b>       | Protein A or G purified from hybridoma culture supernatant                                                                                                                                                                                                                  |
| <b>Immunogen</b>          | <i>E. coli</i> -derived recombinant human XIAP<br>Met1-Ser497<br>Accession # P98170                                                                                                                                                                                         |
| <b>Conjugate</b>          | Alexa Fluor 647<br>Excitation Wavelength: 650 nm<br>Emission Wavelength: 668 nm                                                                                                                                                                                             |
| <b>Formulation</b>        | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide<br><br>*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

|                             |                                                                        |
|-----------------------------|------------------------------------------------------------------------|
| <b>Neutralization</b>       | Optimal dilution of this antibody should be experimentally determined. |
| <b>Immunohistochemistry</b> | Optimal dilution of this antibody should be experimentally determined. |

## PREPARATION AND STORAGE

|                                |                                                                                                                   |
|--------------------------------|-------------------------------------------------------------------------------------------------------------------|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| <b>Stability &amp; Storage</b> | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied                          |

## BACKGROUND

XIAP (X-chromosome linked inhibitor of apoptosis) is a member of the apoptosis (IAP) family of proteins that inhibit caspases. The BIR2 domain of XIAP inhibits caspase-3 and caspase-7. The ability of XIAP to inhibit caspases is prevented by SMAC/Diablo through binding to XIAP-BIR2 and -BIR3 domains.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.