

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human EGFR when phosphorylated at Y1197 in direct ELISAs and Western blots. For Western blot applications, Catalog # MAB8058 is recommended.
Source	Monoclonal Rat IgG _{2B} Clone # 869286
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Phosphopeptide containing the human EGFR Y1197 site Accession # P00533
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

EGF receptor, also known as ErbB1, is an approximately 160 kDa transmembrane receptor tyrosine kinase that binds multiple EGF family proteins. Ligand binding induces EGFR homodimerization or heterodimerization with ErbB2, 3, or 4 as well as activation of its kinase domain and phosphorylation within the cytoplasmic domain. Phosphorylation of Tyr869 by Src is important for full activation of the receptor. Phosphorylation of Tyr1197 by MAP kinases contributes to EGFR interaction with PIK3C2B. EGFR signaling regulates multiple biological functions including cell proliferation, differentiation, motility, and apoptosis. Three additional alternative splice forms lack the transmembrane and cytoplasmic domains. Within the ECD, human EGFR shares 88% aa sequence identity with mouse and rat EGFR.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.